


Porcine and Rangiferine Brucellosis: *Brucella suis*

*Enzootic Abortion,
Contagious Abortion,
Undulant Fever,*

Content Update: June 15, 2007



Importance

Porcine brucellosis, caused by the bacterium *Brucella suis*, is an economically important cause of reproductive losses in pigs. This organism can be maintained in wild and feral swine, complicating eradication efforts in domesticated pigs. One variant, the short rod, is a facultative intracellular pathogen. Other *Brucella* species rarely found in pigs include *Brucella abortus* and *B. melitensis* (For information on *B. abortus* or *B. melitensis*, see the factsheets titled “Bovine Brucellosis” and “Ovine and Caprine Brucellosis,” respectively.)

B. suis contains more diverse isolates than other *Brucella* species, and these isolates have broader host specificity. Five *B. suis* biovars have been identified. Biovars 1, 2 and 3 are maintained in pigs; European hares are also a reservoir for biovar 2. Biovar 4 mainly affects reindeer and caribou and is not normally found in pigs, although is genetically very closely related to biovar 1. Biovar 5 occurs in rodents in the former USSR. Biovar 5 is distinct from other *B. suis* biovars, and may be more closely related to marine mammal *Brucella* isolates.

Some isolates of *B. suis* may be reclassified into a single species (*B. melitensis*), which contains both porcine and marine mammal isolates. This proposal is controversial, and both taxonomic systems are currently in use. Multiple species nomenclature is used in this factsheet.

Species Affected

Most species of *Brucella* are primarily associated with certain hosts; however, infections can also occur in other species, particularly when they are kept in close contact. *B. suis* biovars 1, 2 and 3 affect swine. Biovars 1 and 3 are found in both domesticated pigs (*Sus scrofa domesticus*)

including Siberia, Canada and Alaska. Biovar 5 (murine brucellosis) occurs in the former USSR.

Transmission

In pigs, *B. suis* occurs in the fetus, placenta, fetal fluids and vaginal discharges after an abortion or stillbirth. Pigs usually become infected when they ingest feed contaminated by birth or abortion products, or eat aborted fetuses and membranes. Venereal transmission is also common in swine. *B. suis* is shed in semen; both symptomatic and asymptomatic boars can excrete bacteria.

thrititis, bursitis and osteomyelitis of the vertebral bodies have also been reported.

In hares, *B. suis*

Samples to Collect

***B. suis* biovars 1, 3 and 4 are highly pathogenic for humans; samples should be collected and handled with all appropriate precautions.**

A variety of samples can be collected for culture and microscopic examination. Vaginal swabs, semen or blood samples can be submitted from live animals. Testicles can be submitted after castration. The placenta or aborted/stillborn fetuses can also be cultured. At necropsy, *B. suis* can be isolated from lymph nodes and various organs including the spleen, liver and

Internet Resources

- Centers for Disease Control and Prevention (CDC).
Brucellosis
http://www.cdc.gov/ncidod/dbmd/diseaseinfo/brucellosis_t.htm
- Food and Agriculture Organization of the United Nations.
Manual for the Recognition of Exotic Diseases of Livestock, A Reference Guide for Animal Health Staff
<http://www.spc.int/rahs/>
- Public Health Agency of Canada. Material Safety Data Sheets
<http://www.phac-aspc.gc.ca/msds-ftss/index.html>
- The Merck Manual
<http://www.merck.com/pubs/mmanual/>
- The Merck Veterinary Manual
<http://www.merckvetmanual.com/mvm/index.jsp>
- World Organization for Animal Health (OIE)
<http://www.oie.int>
- OIE Manual of Diagnostic Tests and Vaccines for Terrestrial Animals
http://www.oie.int/eng/normes/mmanual/a_summry.htm

References

- Adams LG, Templeton JW. Genetic resistance to bacterial diseases of animals [online]. Secretariat of Food and Agriculture Organization; 1998. Available at:
http://www.moag.gov.il/brunet/public_sub9_p1.html. Accessed 7 Jun 2007.
- Alton GG, Forsyth JRL. *Brucella* [online]. In Baron S, editor. Medical microbiology. 4th ed. New York: Churchill Livingstone; 1996. Available at:
<http://www.gsbs.utmb.edu/microbook/ch028.htm>. Accessed 4 Jun 2007.
- Centers for Disease Control and Prevention [CDC]. Brucellosis (*Brucella melitensis*, *abortus*, *suis*, and *canis*). CDC; 2005 Oct. Available at:
http://www.cdc.gov/ncidod/dbmd/diseaseinfo/brucellosis_t.htm. Accessed 4 Jun 2007.
- Cutler SJ, Whatmore AM, Commander NJ. Brucellosis--new aspects of an old disease. *J Appl Microbiol*. 2005;98:1270-1281.
- Dieterich RA, Morton JK, Zarnke RL. Experimental *Brucella suis* biovar 4 infection in a moose. *J Wildl Dis*. 1991;27:470-472.
- Gamer G, Saville P, Fediaevsky A. Manual for the recognition of exotic diseases of livestock: A reference guide for animal health staff [online]. Food and Agriculture Or-

- Public Health Agency of Canada. Material Safety Data Sheet – *Brucella* spp. Office of Laboratory Security; 2000 Jan. Available at: <http://www.hc-sc.gc.ca/pphb-dgsp/psp/msds-ftss/msds23e.html>. Accessed 4 Jun 2007.
- Sauret JM, Vilissova N. Human brucellosis. *J Am Board Fam Pract.* 2002;15:401-406.
- Stoffregen WC, Olsen SC, Jack Wheeler C, Bricker BJ, Palmer MV, Jensen AE, Halling SM, Alt DP. Diagnostic characterization of a feral swine herd enzootically infected with *Brucella*. *J Vet Diagn Invest.* 2007;19:227-237.
- U.S. Department of Agriculture, Animal and Plant Health Inspection Service [USDA APHIS]. Wild pigs--hidden danger for farmers and hunters. USDA APHIS; 1992. *Agricultural Information Bulletin* nr. 620. 7 p. Available at: http://www.aphis.usda.gov/lpa/pubs/pub_ahwildpigs.html. Accessed 14 Jun 2007.
- Whatmore AM, Perrett LL, MacMillan AP. Characterisation of the genetic diversity of *Brucella* by multilocus sequencing. *BMC Microbiol* 2007;7:34.
- World Organization for Animal Health [OIE]. Manual of diagnostic tests and vaccines 2004 [online]. Paris: OIE; 2004. Caprine and ovine brucellosis (excluding *B. ovis*). Available at: http://www.oie.int/eng/normes/mmanual/A_00069.htm. Accessed 4 Jun 2007.
- World Organization for Animal Health [OIE]. Manual of diagnostic tests and vaccines 2004 [online]. Paris: OIE; 2004. Porcine brucellosis. Available at: http://www.oie.int/eng/normes/mmanual/A_00096.htm. Accessed 4 Jun 2007.
- World Organization for Animal Health [OIE] Handistatus II [database online]. OIE; 2004. Available at: <http://www.oie.int/hs2/report.asp?lang=en>. Accessed 14 Jun 2007.
- Zygmunt SM, Nettles VF, Shotts EB Jr, Carmen WA, Blackburn BO. Brucellosis in wild swine: a serologic and bacteriologic survey in the southeastern United States and Hawaii. *Am Vet Med Assoc* 1982;181:1285-1287.

*Link defunct as of 2007